


PREPARATION FOR RETRIEVAL – COMMON INFUSION TABLE

USE 50ml LUER LOCK SYRINGES WITH MINIMUM VOLUME TUBING

ADRENALINE **3mg** Dilute to 50ml with 5% dextrose **(60microg/ml)** Commence at 1ml/hr, usual rate 1-20ml/hr



DRAW UP INFUSION DRUG

DRUG	DOSE	DILUTION	CONCENTRATION	TYPICAL RATE
PROPOFOL	500mg	Drawn up undiluted as 50ml	10mg/ml	2-20ml/hr
FENTANYL	500micrograms	Dilute to 50ml with 0.9% NaCl	10microg/ml	5-20ml/hr
MIDAZOLAM	50mg	Dilute to 50ml with 0.9% NaCl	1mg/ml	
KETAMINE	400mg	Dilute to 40ml with 0.9% NaCl	10mg/ml	Loading dose 2-5ml, usual rate 3-20ml/hr
MORPHINE	50mg	Dilute to 50ml with 0.9% NaCl	1mg/ml	
MORPHINE 50mg + MIDAZOLAM 50mg		Dilute to 50ml with 0.9% NaCl	1mg+1mg/ml	Loading dose 2-5ml, usual rate 2-5ml/hr
ADRENALINE	3mg	Dilute to 50ml with 0.9% NaCl	60microg/ml	Commence at 1ml/hr, usual rate 1-20ml/hr
NORADRENALINE	3mg	Dilute to 50ml with 0.9% NaCl	60microg/ml	Commence at 1ml/hr, usual rate 1-20ml/hr

PRIME LINES AFTER MIXING

IF >15mls/hr REQUIRED, DOUBLE INFUSION CONCENTRATION

DRUG	DOSE	DILUTION	CONCENTRATION	RATE
ADRENALINE	6mg	Dilute to 50ml with 0.9% NaCl	120microg/ml	
NORADRENALINE	6mg	Dilute to 50ml with 0.9% NaCl	120microg/ml	

CALL RETRIEVAL CONSULTANT IF DETERIORATION / CHANGED PLAN