

REFERENCES

- Alderman, N. (2007).** Prevalence, characteristics and causes of aggressive behaviour observed within a neurobehavioural rehabilitation service: Predictors and implications for management *Brain Injury* 21(9), 891-911.
- Baguley, I. J., Cooper, J., & Felmingham, K. (2006).** Aggressive behaviour following traumatic brain injury: How common is common? *Journal of Head Trauma Rehabilitation*, 21(1) Jan-Feb 2006, 45-56.
- Eames, P., & Wood, R. (1985).** Rehabilitation after severe brain injury: a follow-up study of a behaviour modification approach. *Journal of Neurology, Neurosurgery and Psychiatry*, 48, 613-619.
- Erg, T. C., Rapport, L. J., Coleman, R. D., & Hanks, R. A. (2002).** Predictors of caregiver and family functioning following traumatic brain injury: social support moderates caregiver distress. *Journal of Head Trauma Rehabilitation*, 17, 155-174.
- Ezrachi, O., Ben-Yishay, Y., Kay, T., Diller, L., & Rattock, J. (1991).** Predicting employment in traumatic brain injury following neuropsychological rehabilitation. *Journal of Head Trauma Rehabilitation*, 6, 71-84.
- Feeney, T., Ylvisaker, M., Rosen, B. H., & Greene, P. (2001).** Community supports for individuals with challenging behavior after brain injury: An analysis of the New York State Behavioral Resource Project. *Journal of Head Trauma Rehabilitation*, 16 (1), 61-75.
- Gardner, R. M., Bird, F. L., Maguire, H., Carreiro, R., & Abenaim, N. (2003).** Intensive positive behaviour supports for adolescents with acquired brain injury: long-term outcomes in community settings. *Journal of Head Trauma Rehabilitation*, 18, 52-74.
- Guercio, J. M. & McMorrow, M. J. (2002).** Proactive protocols for severe unwanted behaviour after acquired brain injury. *Case Manager*, 13, 55-58.
- Hall, K. M., Karzmark, P., Stevens, M., Englander, J., O'Hare, P., & Wright, J. (1994).** Family stressors in traumatic brain injury: a two-year follow-up. *Archives of Physical Medicine and Rehabilitation*, 75, 876-884.
- Johnson, R. & Balleny, H. (1996).** Behaviour problems after brain injury: incidence and need for treatment. *Clinical Rehabilitation*, 10, 173-181.
- Kelly, G., Todd, J., Simpson, G. K., Kremer, P., & Martin, C. (2006).** The Overt Behaviour Scale (OBS): A tool for measuring challenging behaviours following acquired brain injury (ABI) in community settings. *Brain Injury*, 20(3), 307-319.
- Kelly, G., & Winkler, D. (2007).** Long-term accommodation and support for people with higher levels of challenging behaviour. *Brain Impairment*, 8 (3), 262-275.
- Manchester, D., Hodgkinson, A., & Casey, T. (1997b).** Prolonged, severe behavioural disturbance following traumatic brain injury: what can be done? *Brain Injury*, 11, 605-617.
- Marsh, N. V., Kersel, D. A., Havill, J. A., & Sleight, J. W. (2002).** Caregiver burden during the year following severe traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology*, 24, 434-447.
- Rappaport, M., Hall, K. M., Hopkins, K., Belleza, T., & Cope, D. N. (1982).** Disability Rating Scale for severe head trauma: coma to community. *Archives of Physical Medicine and Rehabilitation*, 63(3), 118-123.
- Tate, R. L. (2004).** Assessing support needs for people with traumatic brain injury: the Care and Needs Scale (CANS). *Brain Injury*, 18 (5), 445-460.
- Tate, R. L., Pfaff, A., Veerabangsa, A., & Hodgkinson, A. E. (2004).** Measuring psychosocial recovery after brain injury: change versus competency. *Archives of Physical Medicine and Rehabilitation*, 85(4), 538-545.
- Thomsen, I. V. (1992).** Late psychosocial outcome in severe traumatic brain injury. *Scandinavian Journal of Rehabilitation Medicine*, 26(Suppl), 142-152.
- Walker, A. J., Nott, M. T., Doyle, M., Onus, M., McCarthy, K. (2010).** Effectiveness of a group anger management programme after severe traumatic brain injury. *Brain Injury*, 24 (3), 517-24.
- Watson, C., Rutterford, N. A., Shortland, D., Williamson, N., & Alderman, N. (2001).** Reduction of chronic aggressive behaviour 10 years after brain injury. *Brain Injury*, 15, 1003-1015.
- Wing, J. K., Beevor, A., Curtis, R. H., Park, S. B. G., Hadden, S., & Burns, A. (1998).** Health of the Nation Outcome Scales (HoNOS): Research and development. *British Journal of Psychiatry*, 172, 11-18.
- Ylvisaker, M., Jacobs, H. E., & Feeney, T. (2003).** Positive supports for people who experience behavioural and cognitive disability after brain injury: a review. *Journal of Head Trauma and Rehabilitation*, 18(1), 7-32.
- Zencius, A. H., Wesolowski, M. D., Burke, W. H., & McQuade, P. (1989).** Antecedent control in the treatment of brain-injured clients. *Brain Injury*, 3(2), 199-205.