



**ACI** NSW Agency  
for Clinical  
Innovation

# **ACI Urology Network – Nursing**

## **Male Indwelling Urinary Catheterisation (IUC) - Adult**

### **Clinical Guideline, Competencies & Patient Information Leaflet**

Street address:  
Level 4, Sage Building  
67 Albert Avenue  
Chatswood NSW 2067

Postal address:  
Agency for Clinical  
Innovation  
PO Box 699  
Chatswood NSW 2057

T +61 2 9464 4666  
F +61 2 9464 4728  
info@aci.health.nsw.gov.au  
www.aci.health.nsw.gov.au

# Acknowledgements

Colleen McDonald (CNC Urology: Westmead Hospital), Karina So (CNC Urology: Concord Repatriation General Hospital), Virginia Ip (CNC Urology: Royal Prince Alfred Hospital), Michelle Paul (Nurse Practitioner: Community Health, Greater Newcastle Cluster), Wendy Watts (CNC Urology: John Hunter Hospital) and Suzanne Cruickshanks (CNC Continence: The Continence Service of the University Medical Clinics of Camden & Campbelltown Hospitals). Originally compiled by Urology Nursing Education Working Party Members (September 2008).

**The following pages provide a clinical guideline template to enable clinicians to develop their own resource material relevant to their hospital and Area Health Service. They have been compiled by clinicians for clinicians. If you wish to use this material please acknowledge those that have kindly provided their work to enable use by others. Revise all material with colleagues before using to ensure it is current and reflects best practice.**

**Disclaimer:** The information contained herein is provided in good faith as a public service. The accuracy of any statements made is not guaranteed and it is the responsibility of readers to make their own enquiries as to the accuracy, currency and appropriateness of any information or advice provided. Liability for any act or omission occurring in reliance on this document or for any loss, damage or injury occurring as a consequence of such act or omission is expressly disclaimed.

AGENCY FOR CLINICAL INNOVATION  
Level 4, Sage Building  
67 Albert Avenue  
Chatswood NSW 2067

Agency for Clinical Innovation  
PO Box 699 Chatswood NSW 2057  
T +61 2 9464 4666 | F +61 2 9464 4728  
E [info@aci.nsw.gov.au](mailto:info@aci.nsw.gov.au) | [www.aci.health.nsw.gov.au](http://www.aci.health.nsw.gov.au)

Produced by: ACI Urology Network Nurses Working Group

Further copies of this publication can be obtained from:  
Agency for Clinical Innovation website at: [www.aci.health.nsw.gov.au](http://www.aci.health.nsw.gov.au)

Disclaimer: Content within this publication was accurate at the time of publication. This work is copyright. It may be reproduced in whole or part for study or training purposes subject to the inclusion of an acknowledgment of the source.

It may not be reproduced for commercial usage or sale. Reproduction for purposes other than those indicated above requires written permission from the Agency for Clinical Innovation.

© Agency for Clinical Innovation 2014  
Version: 1.3 March 2017

# TABLE OF CONTENTS

<b>ASSESSING COMPETENCIES</b>	<b>4</b>
What is a competency?	4
<b>COMPETENCY:</b>	<b>4</b>
Male Indwelling Urinary Catheterisation (IUC) - Adult	4
<b>RATIONALE</b>	<b>5</b>
Indications for catheterisation	5
<b>MALE INDWELLING URINARY CATHETERISATION AND CARE - ADULT</b>	<b>7</b>
Definition	7
Expected Outcomes	7
Considerations	7
Equipment	9
Procedure	9
<b>REFERENCES</b>	<b>12</b>
<b>APPENDIX 1.</b>	<b>13</b>
Competency Assessment Forms	13
1. Theory	13
2. Practical	14
<b>APPENDIX 2.</b>	<b>19</b>
Discharge Planning Checklist for Indwelling Catheter	19
Criteria for use of intermittent urinary drainage device – Catheter Valve:	22
<b>APPENDIX 3.</b>	<b>23</b>
Questionnaire	23
Answers	24
<b>APPENDIX 4.</b>	<b>26</b>
Patient Education Leaflet	26
Troubleshooting: Problems You May Encounter	27
<b>APPENDIX 5.</b>	<b>28</b>
Supply of Equipment and Funding Bodies	28

## Assessing Competencies

### What is a competency?

Although the words sound alike, competence and competency are not synonymous. Competence refers to a potential ability, a capability to function in a given situation. Competency is defined as a combination of attributes enabling performance of a range of professional tasks to the appropriate standards. Competency focuses on your actual performance in a situation. This means you need competence before you can expect to achieve competency. Competence enables you to be capable of fulfilling your job responsibilities. Competency however, means that you fulfil your job as expected.

Competency is determined by comparing where you are now with established performance standards developed in the work environment according to your role and setting. Competency standards specify the level of achievement expected and the tasks and contexts of professional practice in which we may see the competency demonstrated.

Competence does not mean expert. Various levels of competence exist and each of these has a minimum acceptable level or standard. Beginners are rarely expert however they can be competent. They perform a wide range of activities methodically and well. The time to complete tasks may take longer however as skill level develops so too will proficiency. They have to ask many questions however are aware of which questions to ask.

### Competency:

#### Male Indwelling Urinary Catheterisation (IUC) - Adult

The aim of the competency is to ensure the Nurse/Medical Officer/ Student is able to demonstrate prior knowledge and can perform male urethral catheterisation to the standard set by the Local Health District or Health Care Provider.

It is necessary that the Nurse/Medical Officer/ Student have theoretical knowledge and technical skill. This knowledge may be gained by attending a study day or undertaking a learning package (discretion of each Local Health District).

The Nurse/Medical Officer/ Student are required to undertake practical sessions with a supervisor in male catheterisation (competency). The supervisor is responsible for identifying the number of supervised sessions for each student in order to be deemed competent. A supervisor must be a Nurse/Medical Officer deemed capable of supervising male catheterisations.

The Nurse/Medical Officer/ Student are responsible for maintaining their skill level. In the event where a Nurse/Medical Officer has been unable to undertake a catheterisation procedure for a significant period and feels they require supervision, it is their responsibility to contact the supervisor for additional supervision. Electronic records will be kept of each Nurse/Medical Officer/ Student that undertakes competency assessment, at the discretion of the Local Health District.

## Rationale

### Indications for catheterisation

- To relieve urinary retention.
- To monitor accurate urinary output.
- To instil medications.
- To manage and maintain urinary system during surgical procedure.
- Establish bladder irrigation for management of haematuria.
- To manage fistula and promote healing.
- To conduct investigative procedures.
- To preserve skin integrity.
- For end-of-life care.

**N.B: Please refer to Local Health District Policy.**

### Competency assessment is required because:

- The ability to insert a catheter safely and effectively on a male patient is an essential clinical skill for Nurse/Medical Officer/Student.
- Competent assessment and education of the patient/client requiring catheterisation is necessary to minimise and prevent complications.

Skill assessment in male catheterisation procedure should be measured against the standard established in the relevant facility policy and protocol. Prerequisites prior to undertaking the skill assessment should include the following:

- Attend in-service or Introduction to Continence Program or a Urinary Catheterisation Workshop.
- Completed online Invasive Device Module (HETI Online).
- Completed online Aseptic Technique Module (HETI Online).
- Completed online Hand Hygiene Module (HETI Online).
- Completed online Waste Management Module (HETI Online).
- Has read the facility specific policy and procedure guidelines on male catheterisation.
- Has knowledge of the clinical indications for insertion of indwelling urinary catheters.
- Observe a Nurse/Midwife/Medical Officer/Student who has achieved competency in this procedure.
- Identify the OH&S principles that apply to the safe insertion of urinary catheter in a male patient.

### Preparation for assessment

- Schedule a time to attend a catheter clinic or liaise with an accredited assessor in your facility.
- Discuss the assessment process with the patient in order to gain verbal consent.
- Community nurses may be able to have their assessments completed in the patient's

home, subject to the patient's consent and following negotiation with the appropriate staff member.

## **Essential components to be assessed**

- Confirm that patient meets Indications for catheterisation.
- Patient gives consent where appropriate.
- Professional attitude and rapport with the patient.
- Comprehensive patient specific assessment and planning.
- Performance of the male patient catheterisation.
- Problem-solving skills and application of theoretical aspect.
- Infection control and aseptic technique.
- Relevant occupational health and safety principles applied to the procedure.
- Patient/carer/family information, education and follow up.

## **Methods of assessment**

- Direct observation of the male catheterisation technique.
- Questioning techniques.
- Demonstrated problem-solving skills in relation to male catheter management.

## **Unsatisfactory Assessment**

In the event of an unsatisfactory assessment the assessor may use the following guide to manage and support staff training:

- The assessor gives feedback on the observed deficits in the performance of the Nurse/Medical Officer/Student.
- The assessor determines with the Nurse/Medical Officer/Student how the skill or knowledge deficit could be best improved.
- Further educational support might be required to prepare the Nurse/Medical Officer/Student for the repeat assessment. This will need to be negotiated between both parties and documented.
- A repeat assessment is to be scheduled.
- If three consecutive attempts are unsuccessful, a further educational, development plan is to be implemented following discussion with the relevant parties. Relevant parties may include the assessor, with the nurse unit manager, nurse educator or clinical nurse consultant.
- Additional education and support are given until competency is achieved.

## **Reassessment/Professional Development**

Reassessment of change of male catheter technique may be required on request.

Nurse/Medical Officer/Student is required to maintain continuous professional development in male catheterisation.

# Male Indwelling Urinary Catheterisation and Care - Adult

## Definition

A urinary catheter is passed through the urethra into the bladder to drain urine. This procedure is performed using sterile equipment under aseptic technique by qualified clinicians.

## Expected Outcomes

1. Urinary catheter inserted only when clinically indicated.
2. The urinary catheter is inserted into the bladder without adverse clinical outcomes.
3. Patient and clinicians safety is maintained.

## Considerations

- Clinical indications for insertion, maintenance and removal of an indwelling catheter must be documented in patient health care records.
- Community nurse may require a Letter of Authority for catheter replacement and removal from the Health Care Team. Please refer to Local Health District Policy.
- Nursing staff should notify Medical Officer of abnormal urine output, less than 30mls in two consecutive hours or urine volume is greater than 1000mls on catheter insertion.
- Long term urethral catheter is usually changed every 4 -6 weeks or as specified by the medical team or certified nurse practitioner / senior urology nurse clinician in some LHDs.
- Catheter urine specimen can be collected from the sampling port within the first 48 hours following catheter insertion. After this period, catheters will be colonized by bacteria. If a urine specimen is required after the first 48 hours, a new catheter should be inserted before the collection is made.
- Patient who is on regular anticholinergic drug for bladder spasms should have the procedure performed one hour after taking the medication.
- If a 2- way urinary catheter is blocked, remove catheter and insert a new catheter and assess possible causes.
- In an acute setting, if patient requires opioid medication before catheter change, the patient has to be monitored for a period of time following the medication guidelines before discharge.
- Another alternative for long term catheters, may need to consider supra pubic catheterisation.

## High Risk for Adverse Outcomes

- **Artificial Heart Valve**
  - Discussion with infection control team
- **Specific Spinal Cord Considerations**
  - If client is a spinal cord injured client above T6 understand autonomic dysreflexia and ensure treatment algorithm for autonomic dysreflexia in spinal cord injury is present
  - If spinal client ascertain if client has used Viagra or Levitra in the last 24hrs or Cialis in the last 4 days as GTN spray, tablet or patches cannot be used

### Alerts

- Do not clamp the catheter in SCI above T6
- Ascertain if client is on anticoagulants prior to procedure
- If the client has an artificial heart valve, discuss antibiotic cover with medical officer prior to change.
- Potential risk of creating a false passage associated with forced instrumentation
- Balloon inflated in urethra/tract resulting in trauma, haemorrhage, rupture or necrosis
- **Autonomic dysreflexia handout to be given to all patients/ clients with a spinal cord injury at or above the 6<sup>th</sup> thoracic level and who have a urethral catheter in situ**

Autonomic Dysreflexia (AD) is a potentially life-threatening condition, which affects people with a spinal cord injury (SCI) at or above the thoracic level 6 (T6). This condition results from widespread reflex activity of the sympathetic nervous system below the level of injury, triggered by an ascending sensory (usually noxious) stimulus. AD can cause a sudden rise in blood pressure that can lead to stroke or even death. One of the common causes is a distended or severely spastic bladder, urinary tract infection, bladder or kidney stones, urological procedure or even inserting a catheter. Episodes of AD could occur during a catheter change. Be alert for sudden hypotension due to rapid draining of bladder or sudden resolution of AD. Initially drain 500mls and then 250mls every 15minutes until bladder is empty. ([http://www.aci.health.nsw.gov.au/\\_data/assets/pdf\\_file/0019/155143/algorithm.pdf](http://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0019/155143/algorithm.pdf))

**Remember: this is a medical emergency and the patient is not to be left alone. Blood pressure should be monitored throughout treatment.**

Patients with a spinal cord injury at T6 and their carers should have a copy of the NSW Health 'Autonomic Dysreflexia Medical Emergency Card' which is available using the following link:

[http://www.aci.health.nsw.gov.au/\\_data/assets/pdf\\_file/0020/163442/Medical-Emergency-Card.pdf](http://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0020/163442/Medical-Emergency-Card.pdf)



## Equipment


- Clean trolley or work surface.
- Adhere to current Hand Hygiene NSW Health Policy and Infection Control NSW Health Policy.
- Catheter pack. This includes
  - Sterile gauze square
  - 10ml Lignocaine 2% sterile lubricant
  - sterile water 10ml ampoule
  - 0.9% Sodium Chloride ampoule 10mls
  - 1pair of sterile gloves
  - 1pair non sterile gloves
  - 2 pairs of forceps
  - Dressing towel
  - fenestrated drape
- 1 sterile Foleys' catheter appropriately sized
- Catheter strap or disposable catheter fixation device
- Sterile drainage bag and or catheter valve
- Protective eye wear
- Disposable protective apron or gown
- Disposable gloves
- Blue disposable sheet
- Rubbish bag
- Specimen jar (if required)
- Sterile kidney dish (if catheter pack has only one tray)
- 30mls of normal saline for irrigation

## Procedure


1. Check correct patient and obtain consent.
2. Perform hand hygiene.
3. If possible, patient should wash their genitals with soap and water or take a shower before procedure.
4. Explain procedure to patient, obtain consent and identify allergies.
5. Ensure patient privacy.
6. Remove underpants and position patient in semi-recumbent position.
7. Perform hand hygiene.
8. Open catheter pack and separate trays, by holding non sterile side of plastic sheeting. Using non touch technique, add equipment to the sterile field.
9. Open sterile gloves.
10. Open sterile drainage bag (if required) and place near patient.
11. Put on apron, goggles and non-sterile gloves.
12. Deflate balloon with 10 ml syringe. Observe amount of water obtained; expect loss of 1ml per week than what was put in.

13. With non-dominant hand hold penis, retract foreskin if necessary and with dominant hand gently rotate and remove catheter.
14. Discard old catheter and perform hand hygiene.
15. Put on sterile gloves.
16. Adjust nozzle of Lignocaine 2% gel and expel air and place on catheter tray.
17. Take one piece of gauze and fold it lengthwise. Separate remaining gauze squares, dip in warm saline and squeeze fairly dry and place in cleaning tray.
18. Open fenestrated drape and place it over patient's genitals.
19. Place cleaning tray just below patient's genital on fenestrated drape.
20. With non-dominant hand, using folded gauze strip hold patient's penis. Retract foreskin if uncircumcised.
21. Using dominant hand, pick up forceps and clean penis with saline swabs from penis tip downwards, one stroke per swab. Discard cleaning tray.
22. Place catheter tray on fenestrated sheet. Holding penis at right angle to the body, insert the Lignocaine nozzle into urethral meatus. Inject the Lignocaine gel into the urethra ensuring firm seal around meatus.
23. Using forefinger and thumb, clamp the urethra for 2-3 minutes, still maintaining penis at right angle to body. Using the Lignocaine syringe massage the under shaft of the penis to assist in moving the gel down the urethra towards the bladder neck or leave catheter syringe in meatus maintaining a seal for 2-3minutes.
24. Pick up catheter with dominant hand, ensuring drainage end of catheter is in tray, gently insert catheter into urethral meatus. When resistance is felt, lower penis and continue till Y junction of catheter. Observe urine flow before inflating balloon, with sterile water (observe patient for any signs of discomfort). Gently withdraw catheter till resistance is felt. Connect catheter to either valve or drainage bag. Observe for urine flow throughout procedure. Do not leave until urine flow occurs.
25. If unable to advance the catheter with firm pressure, abort the procedure and notify Medical Officer
26. Reposition foreskin if required
27. Strap catheter, valve or drainage bag appropriately, ensuring safety, comfort and client's ability to manage catheter.
28. Dispose of waste according to infection control policy.
29. Remove gloves and perform hand hygiene
30. Document on patient integrated notes catheter type, size, balloon size and amount of water in balloon, any abnormalities during the procedure, residual urine colour and volume. Urinary Catheter Record sticker is used in NSW Health public hospitals for easy identification on the Clinical Progress Notes
31. Establish and maintain fluid intake/output chart
32. Also, document order for a catheter review or catheter removal


## Step 1




## Step 2



## Step 3



## Step 4



## References

- Australian Commission on Safety and Quality in Health Care. Standard 3. Preventing and Controlling Healthcare Associated Infections. Safety and Quality Improvement Guide. October 2012.
- European Association of Urology Nurses. Evidence-based Guidelines for Best Practice in Urological Health Care: Catheterisation-Indwelling Catheters in Adults, urethral and suprapubic. 2012. Available from: <http://www.eaun.uroweb.org>.
- Gould CV, Umscheid CA, Agarwal RK, Kuntz G, Pegues DA. and the Healthcare Infection Control Practices Advisory Committee (HIPAC). Guideline for Prevention of Catheter-Associated Urinary Tract Infections 2009. Available from [www.cdc.gov/hicpac/pdf/cauti/cautiguide2009final.pdf](http://www.cdc.gov/hicpac/pdf/cauti/cautiguide2009final.pdf)
- Fekete T. Urinary tract infection associated with urethral catheters. 2013. UpToDate. Accessed January 2014. Available from; [http://www.uptodate.com/contents/urinary-tract-infection-associated-with-urethral-catheters?source=search\\_result&search=urinary+tract+infection&selectedTitle=3%7E150](http://www.uptodate.com/contents/urinary-tract-infection-associated-with-urethral-catheters?source=search_result&search=urinary+tract+infection&selectedTitle=3%7E150)
- Lloyd G, Watts W. A Practice Guide to Urinary Catheterization. Hunter Area Health, NSW; 2003.
- NSW Health (AU) Department Circular, File no.00/7138 Information Bulletin, No.2001/1, Issues 06/2006 on Autonomic Dysreflexia.
- NSW Health (AU). Hand Hygiene Policy – PD2010\_058. Accessed December 2013. Available from: [www0.health.nsw.gov.au/policies/pd/2010/PD2010\\_058.htm](http://www0.health.nsw.gov.au/policies/pd/2010/PD2010_058.htm)
- NSW Health (AU). Infection Control Policy – PD2007\_036. Accessed December 2013. Available from: [www.health.nsw.gov.au/policies/pd/2007/pdf/PD2007\\_036.pdf](http://www.health.nsw.gov.au/policies/pd/2007/pdf/PD2007_036.pdf)
- NSW Health (AU). Medication Handling in NSW Public Health Facilities – PD2013\_043. Accessed 2013. Available from: [www0.health.nsw.gov.au/policies/pd/2013/PD2013\\_043.html](http://www0.health.nsw.gov.au/policies/pd/2013/PD2013_043.html)
- Pomfret I. Catheter care in the community. Nursing Standard. 2000;14(27):46-51.
- Simpson L. Indwelling Urethral Catheterization. Nursing Standard. 2001;15(46):47-53.
- Society of Urologic Nurses and Associates (US) Clinical Practice guidelines, Suprapubic Catheter Replacement 2007.
- The Joanna Briggs Institute (AU) Management of Short Term Indwelling Urethral Catheters to Prevent Urinary Tract Infections, Evidence Based Practice Information Sheet for Health Professionals, 2000. Vol 4 issue 1.

## Appendix 1.

### Competency Assessment Forms

#### 1. Theory

##### Competency: Male Indwelling Urinary Catheterisation (IUC) - Adult

The student is required to show theoretical knowledge of male urethral catheterisation either verbally or in writing prior to undertaking the practical component.

<b>COMPONENTS OF MALE INDWELLING URINARY CATHETERISATION TO BE DISCUSSED</b>	<b>YES</b>	<b>NO</b>	<b>COMMENTS</b>
Described the anatomy and physiology of the urethra, the bladder and abdominal cavity			
Understood the indications for indwelling catheterisation			
Discussed the procedure including the equipment required and the technique			
Discussed types of catheters available, duration of catheterisation, balloon size, closed drainage systems and provided rationale for choice			
Identified complications that can occur during catheterisation and discussed preventative measures and solutions			
Identified problems that can occur during the removal of the catheter and discussed preventative measures and solutions			
Identified care of the indwelling catheter and drainage system			
Discussed special implications related to patients/clients with eg Multiple Sclerosis (MS) and Spinal Cord Injuries at risk of autonomic dysreflexia sexuality, quality of life			
Discussed WH & S considerations			
Discussed legal issues (verbal consent, education, documentation of procedure)			

## 2. Practical

### Changing Male Indwelling Urinary Catheter (IUC) – Adult

Name of Assesse

Signature of Assesse

Ward/Location

Date of Assessment

Name of Assessor

Signature of Assessor

Performance Criteria		
<b>Professional Attitude and Patient Communication.</b>	<b>Yes</b>	<b>No</b>
Introduced self to patient.		
Explained that the procedure for urinary catheterisation is being observed and assessed.		
Gained verbal or inferred consent from the patient.		
Directed visitor / spouse to the waiting area if applicable.		
Addressed any patient concerns that may arise regarding the procedure.		
Performance Criteria		
<b>Patient Assessment and Planning.</b>	<b>Yes</b>	<b>No</b>
Explained the procedure to the patient.		
Stated indication(s) for change of catheter.		
Follow the requirements for patient preparation – as per local policy.		
Provided rationale for catheter selection.		
Identified and planned for potential difficulties.		
Considered the need for pre-procedure medication.		
Performance Criteria on Procedure		
<b>Urethral Catheterisation on a male patient.</b>	<b>Yes</b>	<b>No</b>
Performed hand hygiene.		

Assembled and prepared equipment.		
Positioned patient correctly and continually reassured the patient during the procedure.		
If IUC insitu, put on non-sterile gloves, deflated balloon and removed catheter.		
Put on goggles. Performed hand hygiene and put on sterile gloves. Maintained sterile field and used aseptic technique.		
Cleaned the area using the appropriate solution (as per local policy).		
Correctly placed gauze in 'no-touch' technique.		
Retracted foreskin if present and cleaned glans penis.		
Placed fenestrated drape on patient and while holding penis at 90-° instilled Lignocaine gel. Waited two minutes before inserting the catheter.		
Held penis at a 90° angle and inserted catheter.		
Secured catheter to abdomen.		
Assessed resistance and proceeded cautiously.		
Introduced the catheter to the Y-junction of the catheter.		
Ensured urine flow was present prior to inflating the balloon with sterile water.		
Collected sterile urine specimen if required.		
Connected catheter to the appropriate drainage device.		
Reduced foreskin if retracted earlier.		
Secured catheter and ensured patient comfort.		
Provided appropriate patient education.		
Discarded all equipment appropriately.		
Performed hand hygiene.		
Documented the procedure and any abnormalities.		

Performance Criteria		
<b>Problem Solving Skills. Assessee nurse outlines the reasons for and the management of.</b>	<b>Yes</b>	<b>No</b>
Peri-catheter leakage (bypassing).		
Unable to locate urethral orifice.		
No urine drained on catheter insertion.		
Residual urine is over one litre.		
Performance Criteria		
<b>WH&amp;S Issues Identified and Applied.</b>	<b>Yes</b>	<b>No</b>
Nurse/Medical Officer/Student identifies the following aspects of WH&S and performs a risk assessment prior to performing male urinary catheterisation.		
Positioned patient to minimise need to twist, bend or maintain awkward position. Obtained assistance if required.		
Maintained aseptic technique.		
Used personal protective equipment (facial protection, gown/apron, gloves).		
Correctly disposed of waste.		
Performed hand hygiene in accordance with 5 Moments for Hand Hygiene.		



## Competency: Clamping of Urinary Catheter in a community setting

Clamping of urinary catheter for a short duration, e.g. 30minutes, before catheter replacement procedure, allows the nurse clinician to observe urine flowing freely from the urinary bladder following catheter insertion.

This competency is intended for registered nurses, enrolled nurses and midwives working in community and outpatients settings

Exclusion criteria:

- Clients with spinal cord injury at risk of developing autonomic dysreflexia should not have the urinary catheter clamped before replacement procedure.
- Patient with small bladder capacity, e.g. neurogenic bladder, may not be suitable.

Performance Criteria		
<b>Clamping of Urinary Catheter in a community setting</b>	<b>Yes</b>	<b>No</b>
Educate client or carer on how to place and remove G clamp		
Instruct client to wash hands before handling catheter and G clamp.		
Request client/carer to place G clamp 30 minutes prior to appointment time		
Prepare G clamp by opening space to allow insertion of clamp on tubing of urinary catheter bag.		
Place G clamp on tubing of urinary catheter bag, just below connection of catheter and urine bag.		
<b>Do not place G clamp on catheter as this may affect deflation of balloon.</b>		
Tighten the G clamp to ensure effective occlusion of urine flow.		
Inform client that catheter can be clamped for up to four hours. However if client experience any unexpected physical discomfort related to interrupting urine flow, client should remove G clamp to re-establish free urine drainage system.		
Advise client to wash G clamp in mild detergent after use and store for next use.		
Educate client as per instructions		
Change catheter as per catheter procedure		
Remove G clamp from used catheter bag and return it to client		

## Agreement: Clinical Skill Assessment

### Competency Performance on Male Urethral Catheterisation:

#### Knowledge on male urethral catheterisation procedure

- Satisfactory \_\_\_\_\_
- Unsatisfactory \_\_\_\_\_

#### Male Catheterisation Skill

- Satisfactory \_\_\_\_\_
- Unsatisfactory \_\_\_\_\_

#### Comments by Assessor:

---

---

---

---

---

#### Comments by Assessee (reflection of learning):

---

---

---

---

---

Clinical Assessor (print name): \_\_\_\_\_

Signature \_\_\_\_\_

Date: \_\_\_\_\_

Assessee (print name): \_\_\_\_\_

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

## Appendix 2.

### Discharge Planning Checklist for Indwelling Catheter

#### Urethral catheter

Item	Yes	No	N/A
Educate Patient / Carer / Family on care of the indwelling catheter and drainage system ( including hygiene and problem solving strategies).			
Provide "Catheter care at home" fact sheet.			
Provide minimum one week supply of drainage bags.			
Arrange medical/ Community follow up appointment or Trial of void clinic.			
Provide information on supplier for patient to purchase catheter and drainage equipment.			
Inform patient with permanent catheter on government continence aids assistance schemes.			
Inform patient who is DVA client with gold card of supply arrangement.			
Authorisation letter from Medical Officer for change of catheter by community nurse, if required.			
<p>Name and Signature of Clinician</p> <p>_____</p> <p>Date _____</p>			

## Problem solving with catheters

Problem	Potential Solution
<b>Removal of the indwelling catheter</b>	
Balloon not fully deflated	Refill balloon and allow fluid to enter syringe by gravity (this is called passive balloon deflation).
Cuffing of balloon	Use different type/brand of catheter. Ensure catheter is in the bladder, reinflate balloon with 3 ml of water and allow balloon to deflate passively. Rotate gently whilst removing the catheter.
Bladder spasms	Consider use of anti-cholinergic medication one hour before catheter replacement. Remove catheter slowly as the patient breaths out.
Debris or a stone lodged in the catheter	Drink at least 2 litre of fluid a day whenever possible to reduce the accumulation of debris in the bladder. Use lubricant before applying traction to remove catheter. Refer to a more experienced clinician for advice
Cannot remove catheter	Apply lubricating gel, rotate catheter gently, maintain traction, and ask the patient to bear down. Seek advice from a more experienced clinician when the above options have been tried and problem persists.
Patient anxiety	Reassure client and identify anxiety reasons.
<b>Insertion of catheter</b>	
Difficulty with insertion of urinary catheter	Ensure client is not constipated. Gently rotate the catheter while inserting. Seek advice from a more experienced clinician if unable to advance catheter into the bladder.
Patient anxiety	Reassure client and identify anxiety reasons.
Incorrect catheter size	Check catheter size when preparing equipment. Use the correct catheter size and type. A size 14-16 gauge is usually adequate.

Bladder spasm	Treat symptomatic Urinary Tract Infection (UTI). Take anti-cholinergic medication one hour before the procedure.
Discomfort	Identify cause of discomfort. Ensure adequate lubricant is used. Consider use of anaesthetic gel, wait 3-5 minutes prior to insertion of catheter.
Balloon will not inflate	Remove catheter and insert a new catheter.
<b>Post insertion of catheter</b>	
No urine return	Bladder could be emptied just after catheter removal. Instruct patient to drink a couple of glasses of water before the procedure whenever possible. Do a bladder scan. Patient may have low urine output. Report to Medical Officer.
Catheter bypassing	Correct constipation. Treat bladder spasms. Secure catheter with appropriate device to avoid movement reducing irritation to the bladder. Check catheter patency and drainage bag should be below the level of the bladder. Ensure drainage tubing is not kinked, blocked or looped. If catheter valve is in use, ensure bladder is emptied regularly (frequency can be adjusted according to patient's bladder function e.g. 2- 4 hourly). Ensure catheter size and balloon size is appropriate to patient's condition. e.g. patient has diuresis requires catheter with a bigger lumen. Treat symptomatic UTI.

## **Criteria for use of intermittent urinary drainage device – Catheter Valve:**

- Urine retention.
- Stable detrusor (bladder) function and adequate bladder capacity.
- No recent bladder surgery.
- No suspected bladder perforation.
- No active UTI.
- No active haematuria.
- No vesico-rectal or vesico-colonic tear or fistula.

## Appendix 3.

### Questionnaire

1. What is the preferred balloon size used in a standard Foley Catheter?
2. How often should a leg bag be removed and changed?
3. It is recommended that all silicone and hydrogel coated catheters can be left insitu for how long?
4. What are possible causes for difficult insertion of a male IUC?
5. What action do you take if you experience difficult reinsertion/ removing of an IUC?
6. After insertion of a new catheter what could be the cause of nil return of urine from the catheter?
7. What are the signs and symptoms of autonomic dysreflexia?
8. What action would you take if you attend a patient with autonomic dysreflexia?
9. What issues need to be covered for patient education?
10. What are the principles of closed drainage system?
11. Valves are only suitable for certain patients. Why?
12. How much fluid is recommended for patients with catheter insitu?
13. When would anticholinergics be recommended in a patient with a catheter?
14. How long does it take for Lignocaine to work?
15. What would you document in the notes following a catheter change?
16. What could be the possible reason for difficult inflation of a balloon?
17. What is the accepted hand wash procedure for in the home when changing a catheter?
18. What options exist for the arrangement of supplies for this patient?

## Answers

1. Preferred balloon size used in a standard Foley is 5-10 mls
2. A leg bag should be removed and changed every seven days
3. It is recommended that all silicone and hydrogel coated catheters be left insitu for 4-6 weeks
4. Difficult insertion on a male urethral catheter are possibly
  - Obstructed outlet stricture/prostate
  - False passage
  - Patient anxiety
5. Action to take if difficulty:
  - Removing IDC**
    - Remove excess water from balloon
    - Liaise with expert and leave for another day or person
  - Reinserting IDC**
    - If it is a male try a bigger Foley catheter one size up
    - Stop and inform referring agent for directions
6. Possible causes of nil return of urine post catheterisation are:
  - Catheter not in bladder
  - Catheter lumen blocked with lubricant
7. Signs and symptoms of Autonomic dysreflexia:
  - Rising, high blood pressure
  - Sweating, flushing above level of lesion
  - Pins and needles
  - Pounding headache
8. What to do:
  - Check blood pressure regularly until coming down
  - Remove stimuli: blocked catheter, constipation
  - If still no reduction in symptoms give anti-hypertensive medications
  - If still no reduction in symptoms call ambulance
9. Patient education should include:
  - Hygiene
  - Fluids
  - Care of equipment
  - Signs and symptoms of symptomatic UTI
  - Prevention of constipation
10. Principles of closed drainage are:
  - Sterile system
  - Few disruptions to system as possible
  - Piggy backed night drainage to eliminate disrupting system
  - Change bags weekly and clean overnight drainage bags on daily basis
11. Patients who can have valves must
  - Have normal dexterity or have a carer



- Be motivated
- Have a storing bladder (no urge incontinence)
- No recent bladder surgery
- Have a stable bladder

12. Fluid intake is recommended to be 2 litres unless otherwise indicated by doctor

13. Anticholinergics are recommended when a patient with a catheter is experiencing bladder spasms that are not related to UTI, blockages, or catheter /balloon size but to bladder irritation.

14. Lignocaine works within 2-3 minutes

15. Documentation should include catheter/balloon size, type and material made of, if Lignocaine was used. If any difficulties were encountered with removal or insertion and what sort of drainage was noted. When the next catheter change is due must also be documented. Also document if patient has been experiencing any complication sand what you have suggested they try to address these problems.

16. Difficulty inflating balloon could be

- In wrong spot
- Faulty catheter

17. Hand wash procedure is 2 minute hand wash or use of alcohol rub as per instructions

18. Supply options are:

- Using CAPS entitlement to buy from BrightSky, Independence Australia, Intouch
- Enable NSW
- Private access to individual company

## Appendix 4.

### Patient Education Leaflet

#### Urinary Catheter Care Guide

You have had a urinary catheter inserted through your urethra (water pipe) into your bladder as your bladder is unable to empty by itself. The catheter is held inside your bladder by a balloon, which is filled with water to keep it in place. The catheter is connected to either a urinary drainage bag or a catheter valve.

#### How to Care for Your Catheter

- Wash your hands before you handle your urinary drainage system.
- Empty your urinary drainage bag regularly throughout the day.
- Wash the catheter insertion site everyday with soap and water, and after defaecation.
- Wear supportive underpants to prevent irritation by catheter movement. Urinary drainage bags should be secured firmly but not tightly to the leg using leg straps.
- At night attach the overnight urinary drainage bag to the bottom of the leg bag. Ensure the tap between the leg bag and overnight bag is in the open position.
- Hang the overnight night bag on a hanger, below the level of your bladder. Do not place overnight bag on the floor as this increases the risk of contamination.
- In the morning make sure the outlet tap on the leg bag is closed before disconnecting the overnight bag.

#### Please Ensure That You:

- Drink 2-3 litres of fluid each day unless instructed by your doctor not to. This will promote a steady flow of urine which helps to minimise infection and catheter blockage.
- Avoid constipation as this can prevent the catheter from draining properly.

#### Urinary Drainage Bags

- The overnight bag without a tap is for once only use and should be changed every day.
- The overnight drainage bag with a tap should be cleaned daily and changed weekly.
- The leg bag should be changed weekly.
- The drainage bag should be placed lower than the level of the bladder.

#### Catheter Valve

- Some patients may have a catheter valve instead of a urinary drainage bag.
- The valve must be released every 3-4 hours to empty the bladder (unless uncomfortable, then release as needed).
- Change the valve according to manufacturer's recommendations or Area Health policy. Common practice is to change the valve at the time of urinary catheter replacement.

## **Catheter Change**

- The catheter is changed every 4 - 6 weeks or depending on individual medical condition.
- Your community nurse may instruct you to clamp off the drainage system prior to the catheter change (approx. ½ hour before).
- Ensure that you drink two glasses of water before the nurse is due to arrive.

## **Troubleshooting: Problems You May Encounter**

- It is common to experience bladder irritation or cramps when a new catheter has been inserted. This sensation usually passes in a day or two.

## **If There Has Been No Urine Draining For More Than 2 Hours**

- Check the tubing is not kinked.
- Drink some water - this may flush away any blockage.
- Try walking - this may dislodge a blockage.
- Try milking the catheter or tubing by applying direct pressure.
- If the above fails, contact your nurse or doctor.

## **Contact Your Nurse or Doctor if:**

- Urine is cloudy or strong smelling.
- Urine has become blood stained.
- Pain or tenderness over kidney region.
- Chills or fever.
- Urine is leaking from around the catheter.
- Pain or discomfort from your catheter.

**IF YOU ARE UNABLE TO CONTACT YOUR NURSE OR DOCTOR AFTER HOURS  
PLEASE CONTACT YOUR LOCAL EMERGENCY DEPARTMENT**

## **Contact Details**

**Primary Health Nurse:**

**Family Doctor:**

**Hospital**

## Appendix 5.

### Supply of Equipment and Funding Bodies

An assessment by a continence nurse advisor is recommended to ensure the most appropriate continence product, including the correct fit and application of the product.

#### Enable NSW

Enable NSW provides access to funding for the provision of assistive technology to eligible NSW residents with a permanent or long term disability. Program of Appliances for Disabled People (PADP) is now operated under the umbrella of Enable.

For further information:

Phone: 1800 362 253

Website: [www.enable.health.nsw.gov.au](http://www.enable.health.nsw.gov.au)

### Continence Aids Payment Scheme (CAPS)

This is a federal government scheme available to people aged five years and over who have a permanent and severe incontinence due to:

- Neurological conditions (no Pension Concession Card required) such as intellectual disability (e.g. autism, autism spectrum and Aspergers Syndrome), paraplegia & quadriplegia, acquired neurological conditions (e.g. Alzheimer's Disease, dementia), degenerative neurological diseases (e.g. Parkinson Disease, motor neurone disease), or
- Permanent and severe bladder/bowel innervations (e.g. atonic bladder/hypotonic bladder, prostatectomy with nerve removal) or
- Other causes such as bowel cancer, prostate disease and holds a pension Concession Card

Applicants will need to provide a Health Report from an appropriate health professional such as their medical practitioner or continence nurse about their condition.

Eligible CAPS clients receive an annual indexed payment for continence products

A patient is NOT eligible for CAPS if their incontinence is not permanent or severe or any of the following:

- they are a high care resident in a Australian Government funded aged care home
- they are eligible for assistance with continence aids under the Rehabilitation Appliances Program ( RAP ) which is available through the Department of Veterans' Affairs
- they receive an Australian government funded Extended Aged Care at Home Package (EACH) or an extended Aged Care at Home Dementia Package ( EACH D package )

Further information on eligibility and to obtain an application form:

CAPS Helpline: 1300 366 455

Email: [continence@health.gov.au](mailto:continence@health.gov.au)

## BrightSky Australia offers

- One-stop-shop that provides retail and a national home delivery service of specialist healthcare products.
- Professional continence and wound care advice by phone or appointment. Please call (02) 8741 5600

Address: 6 Holker Street, Newington NSW 2127 (cnr Avenue of Africa)  
Phone no.: 1300 88 66 01  
Fax: 1300 88 66 02  
Email: [orders@brightsky.com.au](mailto:orders@brightsky.com.au)  
Web store: [www.brightsky.com.au](http://www.brightsky.com.au)

## Independence Australia

Independence Australia offers online and retail shopping for medical and healthcare products to the general public. It is also one of the national suppliers of continence products to eligible veterans in Australia under the Rehabilitation Appliances Program (RAP). The order form has to be completed by a health professional.

Showroom Address: 6/ 25-27 Redfern Street, Wetherill Park NSW 2164 (Showroom hours- 9.00am to 4.00pm Monday to Friday)  
Phone: 1300 78 88 55  
Fax: 1300 78 88 11  
Email: [customerservice@independencesolutions.com.au](mailto:customerservice@independencesolutions.com.au)  
Web store: [www.Independenceaustralia.com](http://www.Independenceaustralia.com)

## Intouch Direct

Intouch is one of the national suppliers of continence products to general public, eligible veterans and war widows/widowers.

Phone: 1300 13 42 60  
Fax: 1300 76 62 41  
Email: [healthcare@intouchdirect.com.au](mailto:healthcare@intouchdirect.com.au)  
Web store: [www.intouchdirect.com.au](http://www.intouchdirect.com.au)

## Chemist

You may like to discuss with your chemist about getting your supply and negotiate the price.

## Supermarkets

Incontinence pads are available from local supermarkets.

## **Department of Veterans' Affairs (DVA)**

The Commonwealth Department of Veterans' Affairs (DVA) provides a range of incontinence products to eligible veterans and ward widow/er's via the Rehabilitation Appliances Program (RAP). Eligible applicants need to:

- hold a Gold Card; ( eligible for treatment of all conditions whether or not they are related to war service) ;
- hold a White Card and the incontinence is a result of a specific accepted disability;
- have been assessed by a health professional as requiring products for incontinence;  
or
- products are provided as part of the overall health care management

Gold and White Card holders are not eligible if they are residents receiving high level aged care.

A form requesting the incontinence products is filled out by the assessing doctor or health professional. It is then sent to an authorised product supplier on behalf of the client.

For all enquiries in regards to continence products and supply arrangements, please Contact the South Australian State Office:

National Continence Contract Team  
Department of Veterans' Affairs  
GPO Box 1652  
(199 Grenfell St)  
Adelaide SA 5001  
Phone: 1300 131 945

Or NSW Ministry of Health – Primary Health & Community Partnerships: (02) 9391 9515  
Continence Promotion Centre: (02) 8741 5699